

# **Geophysical and Geological Reconnaissance for the ONR Geoclutter Program**

James A. Austin, Jr., John A. Goff  
University of Texas Institute for Geophysics  
4412 Spicewood Springs Rd., Bldg. 600  
Austin, TX 78759  
phone: (512) 471-0476 fax: (512) 471-0999 email: [goff@ig.utexas.edu](mailto:goff@ig.utexas.edu)

Award Number: N00014-00-1-0844  
<http://www.ig.utexas.edu/research/projects/geoclutter/geoclutter.htm>

## **LONG-TERM GOALS**

The primary goal of the Geoclutter program is to assess geologic clutter/reverberation issues in a seismically and geologically well-characterized shallow-water environment. The mid-outer continental shelf off New Jersey provides such an opportunity, because both bathymetry (a known and prominent cause of backscatter) and portions of the shallow subsurface have been mapped in detail as a result of STRATAFORM. The Geoclutter program consists of three field program phases: (I) an acoustic reconnaissance survey utilizing Navy gray ships and assets to identify potential geoclutter hot spots; (II) a full bistatic acoustic experiment focusing on the chosen areas, and (III), the focus of the work described here, detailed geologic and geophysical surveys of the hot spots identified in Phases I and II.

## **OBJECTIVES**

Our primary objectives for Geoclutter work in 2002 were to (1) continue interpretation of the 2001 and 2002 chirp seismic data, (2) complete analysis of seafloor sample and geotechnical data, and (3) collect and analyze long cores during a test cruise of the AHC-800 drilling system. In the short term, we seek to transition our geologic and geophysical products into reasonable geoacoustic models for use in future acoustic modeling work.

## **APPROACH**


We have employed a variety of approaches in our work. Stratal horizons are interpreted from chirp seismic data using commercial seismic interpretation software. Seafloor measurements, including grain size, porosity, in situ velocity and attenuation, backscatter strength, and acoustic impedance, are compared with each other using correlation analyses. Cores have been both logged for geoacoustic properties, providing ground truth, and sampled, to corroborate geologic interpretation and provide age dating of the sedimentary strata evident in the chirp data.

## **WORK COMPLETED**

Interpretation of the chirp data has continued with significant progress. The analysis of seafloor properties based on in situ acoustic data, grab samples, short cores and remote sensing data (chirp and backscatter) is complete and has been submitted to a peer-reviewed journal (Goff et al., 2003). Three long cores, ranging in length from 4 to 13 m, were collected in 2002 aboard the R/V Knorr using the

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>30 SEP 2003</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2003 to 00-00-2003</b>	
4. TITLE AND SUBTITLE <b>Geophysical and Geological Reconnaissance for the ONR Geoclutter Program</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>University of Texas Institute for Geophysics,,4412 Spicewood Springs Rd., Bldg. 600,,Austin,,TX,78759</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>7</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			


AHC-800 system supplied by DOSECC (Fig. 1). These cores are the longest high-quality cores collected from this part of the margin and represent a unique sample set to provide temporal, stratigraphic and environmental context both for seismic stratigraphic interpretation and future sampling efforts. Geotechnical measurements from these cores were logged at sea, and samples have recently been collected which are currently undergoing detailed analyses for time stratigraphy, sediment texture and paleoenvironmental conditions.



**Figure 1. Location of deep-towed chirp-sonar tracklines collected aboard R/V Endeavor (EN359 and EN370), superimposed on NOAA's bathymetry of the New Jersey middle and outer continental shelf. The small inset locates our study area regionally. Drill sites 1-3 cored with the AHC-800 system aboard R/V Knorr in fall 2002 are marked as yellow stars.**


## RESULTS

Characterization of seafloor samples and *in situ* geotechnical measurements (Goff et al., 2003) illuminate the physical processes of erosion and sediment reworking on the outer shelf that are either active or have been strong seabed modifiers in the recent geologic past. We have also established the relationship between seafloor sedimentology and acoustic backscatter on the New Jersey shelf, and have provided ground-truth for the chirp acoustic impedance measurements with *in situ* velocity measurements (Figure 2).



**Figure 2. Vertical incidence reflection coefficient at 2 kHz versus in situ compressional velocity measurements at 65 kHz.**


Extensive, shallowly buried fluvial/fluviodeltaic/marginal marine dendritic drainage systems have been mapped in Area 1 (Figures 1, 3) using very high resolution (1-4 kHz and 1-15 kHz) geophysical (deep-towed chirp sonar) data. These data record a broad spectrum of channel shapes, depths (from a few to 10's of meters) and widths (~100's to 1000's of meters). We believe that the observed range of channel types is likely influenced by variations in both the magnitude of discharge and the character of the flow regime. A comparison of these mapped systems with a representative group of modern fluvial and marginal marine systems will help us understand the genesis of these shallow subsurface drainage systems.



**Figure 3. A. Interpretation of the dendritic drainage network observed in the Area 1 chirp data. B. Interpreted section of seismic line 26, showing channel fill stratification.**




Preliminary results from the 2001 acoustic reconnaissance experiment have identified Area 2 as a site of interest because of the well-resolved and repeatable clutter targets that were observed there. The Area 2 EN359 chirp data were subsequently augmented during the 2002 EN370 cruise (collected by G.S. Mountain, LDEO and S. Schock, FAU) with ~50 m spacing, for a total of ~750 line km. Interpretations reveal three buried, dendritic and meandering channel systems, ~2-4 km apart and oriented E to NE (Fulthorpe et al., 2002). Channel-fill seismic stratigraphy indicates multiple stages of incision and filling. The northernmost channel system is of primary geoclutter interest. It is mainly V-shaped, up to 350 m wide and 12 m deep, and among the smallest in the Geoclutter survey area. It has been drilled at Site 2, revealing a predominantly sandy fill with, at its base, a burrowed contact with underlying mud that correlates well with the seismic interpretation. A prominent, vertical, seismic facies change, absent from Area 1, occurs throughout Area 2 (Figure 4): stratified material is overlain by up to ~20 m of seismically transparent material separated by a highly irregular boundary, older than the mapped dendritic channel systems. In places, the boundary resembles incisions, which truncate underlying reflections and have flanks that are steeper than even the steepest interpreted channel flanks being mapped in Areas 1 and 2. Partially completed mapping of this facies boundary, using the dense Area 2 profiles, reveals that its irregularities are not entirely random, but the causal mechanism(s) for these features is/are as yet unknown.



**Figure 4.** Underway interpretation of the facies change horizon being mapped in Area 2.

Examination of the geologic and climatic history of a continental margin using seismic reflection data is dependent on identifying key horizons that mark geologically and climatically significant episodes. On the New Jersey margin, the “R” horizon has been postulated to be a key indicator of the climate just prior to the end of the Last Glacial Maximum (LGM), possibly the major erosional event that immediately preceded the deposition of post-LGM sediments. Post-LGM sediments have been subsequently eroded by a series of fluvial channels that appear to correlate in places with acoustic clutter. The suite of ultra-high resolution chirp-sonar reflection data which clearly image “R,” and Site 3 of the 2002 drilling cruise on the *R/V Knorr* which penetrated “R”, are excellent tools for investigating the geographic distribution and sequence stratigraphic relationship for the “R” reflector. A preliminary interpretation of “R” within the Area 1 and along Area 3 dip lines has been completed (Figure 5).



*Figure 5. Interpreted and interpolated “R” horizon in Area 1 and along dip profiles.*

## IMPACT/APPLICATIONS

The primary application of our geological and geophysical characterization is in the establishment of critical paleoenvironmental characteristics for the understanding of acoustic interactions with the seabed.

## RELATED PROJECTS

The ONR STRATAFORM program provided initial site characterization for the Geoclutter natural laboratory. The SWAT acoustic experiment was also carried out in this area, and the 2006 Shallow Water Acoustics experiment is now planned for this area.

## REFERENCES

- Goff, J. A., B. J. Kraft, L. A. Mayer, S. G. Schock, C. K. Sommerfield, H. C. Olson, S. P. S. Gulick, and S. Nordfjord, Seabed characterization on the New Jersey middle and outer shelf: Correlability and spatial variability of seafloor sediment properties, submitted to *Mar. Geol.*, Aug. 2003.
- Fulthorpe, C. S., J. A. Goff, J. A. Austin Jr., S. P. Gulick, and S. Nordfjord, Late Quaternary Incisions and Related Shallow Subsurface Stratigraphy on the New Jersey Mid-Outer Shelf: Preliminary Results from Ultra-High Resolution Chirp Sonar Images - Part II, *Eos Trans. AGU*, 83, Fall Meet. Suppl., Abstract OS71C-0300, 2002